

胸壁、肋膜及縱膈腔病變

三軍總醫院

胸腔內科暨睡眠中心主任

彭忠衍醫師

病灶的定位與辨別

■ 先判斷胸腔內或胸腔外 → lateral view、PE

■ 胸腔外 (extra-thoracic)

- Skin

- Foreign body

■ 胸腔內 (intra-thoracic)

- 肺內 (intrapulmonary) : 肺實質病灶

- 肺外 (extrapulmonary) : pleura, chest wall (bone, soft tissue)

Intrapulmonary vs. Extrapulmonary

(肺内)

(肺外)

- **Incomplete border sign**
- **Tapered margin sign**
- **Center outside the lung**
- **Bilateral lesion**

Imaging of Chest Wall Lesion

Chest Wall Lesions – 軟性胸壁

■ Foreign body

- button, electrode, wires, tube, dressings and hair braids (髮辮)

■ Skin tumor

- Neurofibromatosis, moles / melanoma, Kaposi's sarcoma

■ Soft tissue tumor

- Fibroma, lipoma, hemangioma, muscle tumor

■ Calcification

- Parasite calcification (Filaria cysticerci)
- Granulomatous LAP

■ Subcutaneous emphysema

- Pneumothorax: trauma, post-procedure....
- Deep neck infection

■ Breasts / nipples

- s/p mastectomy
- Breast tumor
- Mammoplasty

Breast

■ 正常的乳房影像：

- 為一均質的陰影，由上而下直到橫膈，由內而外直到胸壁，其濃度會逐漸增加。如果這個陰影沒有造成lung marking的減少或遮蔽，則可以判定是來自乳房造成的陰影，而不是肺內病變。

■ 異常的乳房影像：

- Mammoplasty：沒有正常乳房的濃度漸增現象
- 單側乳房異常：s/p mastectomy, breast tumor

Chest Wall Lesions – 骨性胸壁

■ Sternum:

- Funnel chest (pectus excavatum, 漏斗胸)
- Pigeon chest (pectus carinatum, 鸽胸)

■ Spine:

- Kyphoscoliosis
- Neurogenic lesions
- Compression fracture
- Osteopenic / osteogenic lesion of metastasis
- Paraspinal abscess

Chest Wall Lesions – 骨性胸壁

■ Rib:

- Costal cartilage: 分辨性別、年齡
- 先天發育異常: forked rib, fusion rib, cervical rib
- Contour異常: rib fracture; funnel chest, barrel chest
- Rib notching (下緣): CoA, neurogenic tumor (benign)
- Rib變寬/size變大: thalassemia, bone tumor
- Density改變:
 - ↑(變白): diffuse - **2M** (metastasis, myelofibrosis), **2O** (osteopetrosis, osteodystrophy); 鈣化 - Osteochondroma, chondrosarcoma, osteosarcoma
 - ↓(變黑): osteolytic, osteoporosis
- Chest wall mass + rib destruction:
 - 成人: Multiple myeloma (plasmacytoma), metastasis, Pancoast tumor
 - 小孩: Ewing's sarcoma, neuroblastoma
 - 感染: TB, aspergillosis, actinomycosis, nocardiosis

用Rib來判斷病患的性別和年齡

■ 性別：calcification of costal cartilage的形態

■ M: peripheral (vaginal type)

■ F : central (penial type)


■ 年齡：

■ 第一根肋軟骨鈣化: 30-40 y/o

先天發育異常

■ Bifid or forked rib

- 最常發生於4th rib
- The anterior portion of rib is duplicated



先天發育異常

■ Cervical rib:

- A supernumerary or accessory rib arising from the **C7** vertebra.
- About 0.5% of the population and is more common in **females** than in males.
- Usually asymptomatic
- May **thoracic outlet syndrome** by compression of the brachial plexus or subclavian vessels.
 - Pain in the hand when the arm is elevated
 - Difference in pulse intensity between the two arms when the affected extremity is in a certain position

Rib Notching (下緣)

Coarctation of Aorta-主動脈狹窄

- Deformity of the aortic media and intima → prominent posterior infolding of the aortic lumen.
- Characteristically occurs at or near the junction of the aortic arch and the descending thoracic aorta
- Two main types:
 - **Localized (postductal or adult-type):** most common
 - a focal narrowing of the aorta, just beyond the origin of the left subclavian artery or the ligamentum arteriosum
 - accompanied by dilatation of the left subclavian artery
 - **Tubular hypoplasia (preductal or infantile-type)**
 - The 2nd common cause of heart failure in newborn
 - A long narrowed segment beyond origin of innominate artery
 - Associated with intracardiac defect, esp. a deformed or bicuspid aortic valve

Rib Notching (下緣)

Coarctation of Aorta-主動脈狹窄

■ Imaging

■ Inferior rib notching

■ Best diagnostic clue

- An elaborate system of **collateral** vessels (including collateral internal mammary, intercostal, and superior epigastric arteries) forms to bypass the coarctation.
- The dilated and tortuous intercostal vessels form deep grooves on the undersurfaces of the ribs
- Usually 3rd ~8th ribs
- 1st and 2nd ribs supplied from costocervical trunk, not from D-aorta, so both not served as collateral vessels

Table 1
Radiologic Differentiation of Malignant Chest Wall Tumors

Imaging Finding	Tumor Type
Fat component	Liposarcoma
Calcification	
Skeletal	
Rings and arcs	Chondrosarcoma
Flocculent or stippled	Chondrosarcoma
Centrally dense	Osteosarcoma
Extraskeletal	
Heterogeneous	Ganglioneuroblastoma or neuroblastoma
Speckled	Proximal-type epithelioid sarcoma
Diffuse osteolytic change	Myeloma
Ill-defined mass	
Eccentric growth, in children and young adults	Ewing sarcoma
Fluid-fluid levels and calcification, in adolescents and adults	Synovial sarcoma
Chronic lymphedema	Angiosarcoma
Infiltrative growth	Malignant lymphoma
Nonspecific findings	Leiomyosarcoma, rhabdomyosarcoma, malignant fibrous histiocyoma, aggressive fibromatosis, malignant peripheral nerve sheath tumor, or dermatofibrosarcoma protuberans

Imaging of Pleural Lesion

Pleural Lesions

- **Pleural effusion**

- **Pleural mass**

- **Pleural thickening / calcification**

- **Pneumothorax**

Pleural Effusion

- **Free pleural effusion**
- **Subpulmonic effusion**
- **Encapsulated pleural effusion**
- **Interlobar pleural effusion**

Free Pleural Effusion

- Normally, 10-15cc in one pleural space
- 隨姿勢改變形狀
- PA view:
 - Lateral C-P angle or medial phrenicovertebral sulcus
 - Meniscus sign or angle blunting
 - 通常積水量 > 300cc
- Lat view:
 - Posterior CP angle blunting，積水量 > 150cc
 - More sensitive than PA view
- Supine view:
 - 由橫膈至肺尖呈現homogenous GGO (略有漸層)
 - 當lateral CP angle blunting時，積水量約500~1000cc
- Decubitus view:
 - Can detect PE less than 100cc



Meniscus sign

- Pleural effusion為底盤寬大、上方窄細的形狀


- 與重力有關

- Meniscus sign的形成

- 外側的lung厚度較薄，趨近中央時肺的厚度逐漸增加 – Density titration by air
- 在平行X-ray beam的方向上，在periphery處X-ray beam需要穿過greater depth of effusion ($e > a-a'$)，所以愈內側的superior margin of pleural effusion在CXR上愈不容易看出來



Decubitus position for D/D of pleural effusion or thickening



Subpulmonary Pleural Effusion



■ 定義

- Pleural effusion 只會積在下肺葉和橫膈之間，而不會溢出至同側的C-P sulcus。
- 由於積液積存於橫膈和visceral pleural of lower lobe之間，積液量愈多，則lower lobe愈往上推移，所以CXR上有如橫膈一般，稱為“pseudodiaphragm”。

■ PA view

- Diaphragm 的最高點向外移至外 1/3 (正常的在內1/2~1/3)
- C-P angle變鈍、變淺
- Diaphragm以下的lung markings 消失
- L't side: diaphragm 和 gastric bubble 相隔>2cm

D.D. of Pleural Effusion (1)

■ With cardiomegaly:

■ H (hemodynamic)

■ Cardiovascular:

- CHF

- Post-cardiotomy syndrome

■ Collagen-vascular disease

- SLE, RA

■ Pulmonary embolism with R't side heart enlargement

■ I (infection): myocarditis or pericarditis with pleurisy

- TB, virus, rheumatic fever

■ N (neoplasm): malignant pericardial effusion and pleural effusion

- Metastasis, mesothelioma

■ With subsegmental atelectasis:

- Post-operative

- Pulmonary embolism

- Rib fractures

- Abdominal mass / abscess / ascites

D.D. of Pleural Effusion (2)

■ With lobar opacities:

- Pneumonia with PPE/
empyema
- Chronic infection :
 - TB (common)
 - Fungus (rare)
- Pulmonary embolism
- Neoplasm:
 - Lung ca
 - Lymphoma
 - Metastasis

■ With hilar enlargement:

- Vascular shadow
 - Pulmonary embolism
- Lymphadenopathy
 - Neoplasm
 - Lung ca.
 - Metastasis
 - Lymphoma
 - Infectious :
 - TB
 - Fungus infection (rare)
 - Inflammatory:
 - Sarcoidosis

Pleural Masses

■ Solitary

- Loculated fluid
 - 水 (pleural effusion)
 - 血 (hemothorax)
 - 膿 (empyema)
- Neoplasms
 - Mesothelioma
 - Metastasis
 - Fibrous tumor of pleura
- Mesothelial cyst

■ Multiple

- Loculated pleural effusion
- Neoplasms (3M)
 - Metastases (Adenoca)
 - Malignant mesothelioma
 - Malignant thymoma
- Pleural plaque

Pleural Metastasis

■ Origin:

- 40% from lung carcinoma
- 20% from breast carcinoma
- 10% from lymphoma
- Remaining 30% from other primary sites

■ Image:

- May mimic malignant mesothelioma
- Malignant effusion is the most frequent manifestation of metastatic pleural disease and is often accompanied by solid tumor deposits of variable size

Mesothelioma

- 又叫做Localized fibrous tumor of pleura
- Usually benign; 37% malignant change
- Mesothelioma
 - Benign (non-cancerous) mesothelioma
 - Solitary fibrous tumor
 - 與asbestos無關
 - 主要在肺的周邊或是位於fissure
 - Malignant mesothelioma
 - Solitary or multiple
 - 與asbestos有關
 - Invasive and destructive

Malignant Mesothelioma

- The most common primary neoplasm of the pleura
- Risk factors:
 - Asbestos, radiation, chronic pleural disease
 - Not related to cigarette smoking
- Asbestos pleural plaque is not the precursor of mesothelioma
- Latency from the first exposure to asbestos: 35~40 yrs
 - Peak in 60~70 y/o
 - Rare in childhood
- **Male** predominant (4x)
- Poor prognosis, most p't dying within 1 yr

Malignant Mesothelioma

■ Image:

■ Extensive, lobulated nodular pleural thickening

- CT scan: tumor thicker at base
- The most common findings on CT scan

■ Pleural effusion

- **Unilateral**, rarely bilateral (10%)
- Moderate to large amount (1/3高度: 50%, 2/3高度: 40%, > 2/3高度: 10%)

■ Small hemithorax (contracted)

- 有時會因為pleural effusion而造成volume expansion (15%)，但是很少造成明顯的mediastinal shift

■ Other findings of CT scan

- Mediastinal organ invasion: 可能包圍mediastinal structure > 50%
- Pleural plaque over contralateral lung
- Transdiaphragmatic extension

Differential Diagnosis of Benign and Malignant Pleural Disease

■ Favor malignant pleural disease

■ Circumferential involvement

■ Nodular thickening

■ Parietal pleural > 1cm

■ Mediastinal pleural involvement

Differential Diagnosis of 3”M”

- Malignant mesothelioma

- Pleural effusion (90%)
- Extensive lobulated pleural thickening
- Often contracted hemithorax

- Malignant (invasive) thymoma

- Anterior mediastinal mass

- Metastasis of pleura (esp. adenocarcinoma)

- Less often contracted hemithorax

Pleural Thickening

■ H (hemodynamic)

- Healed (organized) hemothorax
- Collagen-vascular disease: 因反覆pleural effusion

■ I (infectious)

- Organized empyema related to pyogenic or TB infection:
最常見

■ I (inhalational)

- Asbestosis, talcosis

■ N (neoplasm)

- Diffuse nodular pleural thickening
 - 3M + primary lung cancer (adenocarcinoma)
- Pancoast's tumor:
 - Apical pleural thickening + bony destruction

Pleural Calcification

■ 原因：

- Fibrocalcified visceral pleura:

 - Pyothorax, TB

 - Hemothorax

- Fibrocalcified parietal pleura:

 - Asbestos

 - Talc

- 兩側橫膈和lateral chest wall的pleural calcification: **asbestosis** or **talcosis**

Pneumothorax

■ Upright patient:

- Identify the **pleural line**
- Most often near the **apex**

■ Supine patient:

- 在supine position，胸腔最高處為**anterior C-P sulcus**，其次為**lateral CP sulcus**。所以發生氣胸時，air最先填充至anterior CP sulcus，量更多時，lateteral CP sulcus亦被填滿。
- **Deep sulcus sign**: anterior和lateral CP sulcus顯得非常深，橫膈被襯托得很清晰
- Sharper than normal appearance of hemidiaphragm

■ When pleural line is not identified:

- Lateral decubitus view
- Expiratory film: no more useful

Imaging of Diaphragm & Sub-diaphragmatic Lesions

Diaphragm Depression

■ Bilateral

- Emphysema
- Bilateral pneumothorax
- 瘦長體格

■ Unilateral

- Tension pneumothorax
- 下肺野有大的bullae
- 單側lung hyperexpansion
(ex: 吸入異物)

Diaphragm Elevation

■ Bilateral

■ 肺部因素

- 吸氣不足or 吐氣照相
- Bilateral atelectasis
- Restrictive lung disease

■ 腹部因素

- 油：obesity
- 水：ascites
- 小孩：懷孕
- 器官：肝脾腫大

■ Unilateral

■ 肺部病灶

- Subpulmonary effusion
- Decreased lung volume

■ 腹部病灶

- Abdomen mass: liver, distended stomach, colon interposition
- Subphrenic abscess

■ 橫膈本身病灶

- Eventration
- Diaphragmatic hernia
- Traumatic rupture of diaphragm
- Diaphragmatic tumor
- Phrenic nerve paralysis

Decreased Lung Volume

- Atelectasis / collapse
- Lobectomy / pneumonectomy
- Hypoplastic lung

Abdominal Diseases

- Subphrenic abscess

- Elevated diaphragm + pleural effusion 最常見
- Localized extraluminal air below the diaphragm
- Basilar platelike subsegmental atelectasis

- Liver:

- HCC
- Liver abscess: air-fluid level within liver shadow

- Distended stomach

- Interposition of colon

Diaphragm Hernia (1)

■ Hiatal hernia

- The most common non-traumatic hernia
- Risk factors: obesity, pregnancy
- A retrocardiac mass, usually with air or air-fluid level
- Near the midline or cross the it
- Occasionally on one side and may mimic lung abscess

Diaphragm Hernia (2)

■ Morgagni's hernia

■ 右前、内

- More common in adult than in infant
- Risk factors: obesity, effort, increased abdominal pressure
- Herniated abdominal contents (in order of decreased frequency): omentum, colon, stomach, liver, and small intestine
- CXR: a **smooth, well-defined opacity** in **right cardiophrenic angle**, may obscure the right heart border.


Diaphragm Hernia (3)

■ Bochdalek's hernia

- Usually diagnosed in infants who present with clinical symptoms of pulmonary insufficiency.
- A lung-based soft-tissue-opacity lesion; left (70%–90%, presumably owing to the protective effects of the liver), posterior (左後)
- Must be considered when the mass is continuous with the diaphragm
- The herniated contents mostly contain fat and omental tissue.



赫尼亞
Bochdalek's hernia



Axial hiatal hernia
軸狀裂口赫尼亞



食道赫尼亞
Paraesophageal hernia




Morgagni 赫尼亞



Mediastinum Lesions

Anatomy Landmarks

- Identify the location to narrow D/D
 - PA view: Silhouette sign
 - Lat. View: better
- Divide anterior and middle mediastinum:
 - Anterior border of the trachea – posterior border of the heart – posterior border of IVC
- Divide middle and posterior mediastinum:
 - A line 1cm posterior to the anterior border of the vertebral bodies



- **Anatomy Landmarks**

- 氣管前緣、心臟後緣、IVC
- 椎體前緣向後 1 cm 畫線

- 分為前、中、後縱隔

A line, 1cm back from the anterior edge of the vertebral bodies

List of Mediastinum Lesions

- Shift

- Widening

- Aortic aneurysm / dissection
- Mediastinitis ($\pm$ air-fluid level)
- Lipomatosis

- Soft tissue density

- Mass, Neoplasm

- Air or air-fluid level

- Pneumomediastinum
- 食道：
 - Esophagus reconstruction
 - Achalasia
- Hiatal hernia

Mediastinal Shift

- Mechanism: Pressure imbalance
- Determine which side is diseased by
 - Associated findings
 - Expiratory film: air-trapping
 - Lateral decubitus film
- Causes
 - Lung volume:
 - Decrease (ex: atelectasis, pos-OP),
 - Increase (ex: check valve; air進得去，但出不來)
 - Huge lung mass
 - Pleural disease (effusion, air, mass)

List of Mediastinum Lesions

- Shift

- Widening

- Aortic aneurysm / dissection
- Mediastinitis ($\pm$ air-fluid level)
- Lipomatosis

- Soft tissue density

- Mass, Neoplasm

- Air or air-fluid level

- Pneumomediastinum
- 食道：
 - Esophagus reconstruction
 - Achalasia
- Hiatal hernia

Mediastinal Widening

■ First, exclude below conditions

- Supine AP film
- Lordotic film
- Inadequate inspiration

Mediastinal Widening

■ Hemodynamic (Vascular)

- Tortuous aorta, aortic aneurysm, aortic dissection
- Hematoma: trauma, post-operation, catheters

■ Mediastinitis

- Acute mediastinitis
- Fibrosing mediastinitis

■ Neoplasm / Mass

- Lymphoma, lung ca, LAP

■ Lipomatosis

- Steroid use, Cushing's syndrome, obesity

Acute Mediastinitis

■ Causes:

- Esophageal perforation: the most common cause (Boehaave's syndrome, TE fistula)
- Esophageal/cardiac surgery
- Direct extension of infection from adjacent structures (e.g. retropharyngeal abscess)
- Hematogenous spread of infection

■ CXR:

- Widening of mediastinum, typically with blurred ill-defined mediastinal margin
- Pneumomediastinum may be evident

■ Diagnosis:

- By extravasation of ingested contrast into the mediastinum or pleural space

Pneumomediastinum

- Gas within the mediastinum can result from 5 sites:
 - Lung (most common), mediastinal airway, neck, esophagus, and abdominal cavity.
- CXR:
 - Lucent streaks of gas outlining the mediastinal structures
 - **Continuous diaphragm sign**: when gas interposed between the heart and diaphragm, it permits identification of the central portion of diaphragm.

Pneumomediastinum

- The terminal air spaces of the lung are the most common source of pneumomediastinum.
- In the presence of a pressure gradient between an alveolus and the interstitium, air ruptures from the alveolus into the perivascular and peribronchial fascial sheath (interstitial emphysema)
- Continued insufflation caused an overflow of air into the retroperitoneum, anterior mediastinum, and subcutaneous tissues of the neck and axilla

List of Mediastinum Lesions

- **Shift**
- **Widening**
 - **Aortic aneurysm**
 - **Lipomatosis**
 - **Mediastinitis (air-fluid level)**
- **Soft tissue density**
 - **Mass, Neoplasm**
- **Air or air-fluid level**
 - **Pneumomediastinum**
 - **Esophagus reconstruction**
 - **Achalasia**
 - **Hernia**

Hint of Mediastinal Mass


- Tapered border
- Center within mediastinum: 圓心落在縱膈腔內
- Sharp margin
- Bilaterality: 病灶跨在左右兩側的胸腔

Anterior mediastinum

Substernal thyroid gland




Thymoma



Teratoma

Lymph nodes; lymphoma

Tumors of Mediastinum



Middle mediastinum

Vascular; aneurysm, enlarged heart

Lymph nodes; lymphoma, metastatic cancer

Esophageal; achalasia, diverticula


Bronchogenic or pericardial cyst

Posterior mediastinum





Neurilemoma
Neurofibroma
Ganglioneuroma
Schwann-cell tumor






Extra-pulmonary sign



Intra-pulmonary lesion



Mediastinal Compartments


Anterior mediastinal mass

1. Pre-vascular : 3T + 1L

2. Cardiophrenic space: 4種D/D

- heart
- mediastinum
- lung
- diaphragm

- Anterior
- Middle
- Posterior



判讀 Anterior Mediastinal Mass 的一般原則

■ 定位：

- Mediastinal (junction) line: Ant. disrupted, Post. preserved
- Hilum overlay sign(+): anterior or posterior to hilum
- Location of lesion: 與diaphragm接觸的lesion，如pericardial cyst/aneurysm, Morgagni hernia, pleuropericardial cyst
- 利用anatomy的silhouette sign:
 - A-aorta, SVC, heart border: anterior
 - Aortic arch: middle
 - D-aorta: posterior

■ Tissue density

Thymic Lesions

- Thymoma :

- 以是否有 **local invasion at surgery** 來分類

- 分為invasive or non-invasive

- 比較不prefer benign or malignant的分類

- Thymic carcinoma

- Thymic cyst

Thymoma(1)

- **Normal thymus:** located in the superior anterior mediastinum, usually anterior to the proximal A-aorta and distal SVC.
- 最常見的原發性前縱膈腔腫瘤，約20%
- Middle-aged male adults(> 40 y/o)
- 10~15% MG p't有thymoma，35~40% thymoma p't有MG
- Pure RBC aplasia (PRCA): thymoma最常見的 hematologic abnormality, 5%
- Slow-growing, encapsulated
- Recurrence or local invasion常發生於anterior mediastinal soft tissue, pericardium, or pleura

Thymoma(2)

■ CXR:

- Most near the junction of the heart and great vessels
- Round or oval; smooth or lobulated margin
- Anterior sulcus sign: when touching the sternum, not flattened against it (very firm)
- Protrude to one or both sides of mediastinum
- Calcification: thin, linear and peripheral or throughout its substance
- Radiographic appearance between **invasive** or **non-invasive** thymoma is **indistinguishable**.

★ ■ **Invasive thymoma**: rarely metastasis, but more commonly extends locally or seeds the pleural or pericardial surfaces.

■ CT:

★ ■ **Bilaterality, larger size, lobulated contour, poorly-defined margin, and associated pleural effusion/nodules/masses** – favor **invasive thymoma**

Thymic Cancer

- CXR: lobulated irregular marginated large anterior mediastinal mass
- CT:
 - **Heterogeneous enhancement** with area of necrosis
 - Usually aggressive invasion of pericardium and pleura, great vessels and other mediastinal structures

Thymic Cyst

- CXR: well-marginated anterior mediastinal mass
- CT:
 - Anterior mediastinal mass with a single dominant or multiple thin-walled cysts

Germ Cell Tumor

■ Classifications:

- Teratoma

- Seminoma

- Non-seminomatous germ cell tumor (NSGCT)

Teratoma

- Most common mediastinal GCT; 70%
- Most often in children or young adult(20-30y/o); 男=女
- 3 histologic types:
 - Mature teratoma: benign
 - Immature teratoma: low potential to malignancy
 - Teratoma with malignant transformation: aggressive malignant
- CXR:
 - Lopsided(偏向一邊), lobulated, well-marginated mass; most common in anterior mediastinum(85%), usually near thymus
 - 25% calcification
 - Soft, tend to be flattened against mediastinum: 與thymoma不同
- CT:
 - Multiloculated cystic mass with variable thickness wall(80%), may contain fat, fluid, soft tissue and calcification densities; finding of teeth is pathognomonic

Calcified Mediastinal Mass

■ 3T+1L

- Teratoma
- Thyroid goiter
- Thymoma
- “Treated” lymphoma

■ Calcified LAP

- Fibrosing
mediastinitis:
histoplasmosis, TB
- Silicosis
- Sarcoidosis

■ Aortic aneurysm

■ Rim calcification

- Cystic teratoma
- Thymic cyst
- Aortic aneurysm
- Duplication cyst
 - Bronchogenic cyst
 - Neuroenteric cyst

Thyroid Lesions

- The most frequent pathologic finding is **multi-nodular goiter**
- 75-80% arise **from a lower pole or the isthmus** and extend into the anterior or middle mediastinum.
- Frequently associates with a palpable **neck mass**
- D/D with other thoracic inlet lesions

Thyroid Lesions

■ CXR:

- ★ ■ 邊緣平滑清楚的mass + tracheal deviation at the level of thoracic inlet.
- ★ ■ Tend to extend above the clavicle.
 - 25% calcification
 - Lat. view: anterior goiter fill in **retrosternal clear space**, and **deviate trachea posteriorly**.

■ Chest CT:

■ NECT:

- High attenuation due to natural iodine
- Sharply demarcated heterogeneous mass

■ CECT:

- Strong enhancement

Lymphoma

- As a mediastinal tumor, **H.D.(85%) > NHL(50%)**
- CXR:
 - Most commonly involve anterior, superior mediastinal LN (prevascular, paratracheal), bilateral, bulky and asymmetric
 - Bilateral mediastinal widening with lobulation
 - 20% calcification after R/T; rim or mulberry(桑椹)
 - Lung involvement: usually **peripheral** and **subpleural**
 - Consolidation with air-bronchogram
- **v** ■ **Single or multiple discrete pulmonary nodules/masses, less well-defined and less dense**
 - Diffuse reticulonodular opacities (LIP)
- Chest CT
 - Minimally enhanced in CECT
 - 通常會 **encase or displace** 鄰近的 structures，很少會侵犯或是造成狹窄。

Middle Mediastinum

Table 2
Middle Mediastinal Masses

Lymphadenopathy

Aortic arch aneurysm

Enlarged pulmonary artery

Foregut duplication cysts (bronchogenic, esophageal, neurenteric)

Pericardial cyst


Tracheal lesions

Mediastinal Compartments

Middle Mediastinal Lesions

1. LAP
2. Vascular lesions
3. Esophageal lesions
4. Tracheal lesions
5. Duplication cysts

- Anterior
- Middle
- Posterior



Lymphadenopathy (LAP)

- Neoplastic
- Infection / Inflammatory
- Inhalational

Neoplastic Adenopathy

- **Neoplastic LAP** is the most common middle mediastinal lesions
- Causes :
 - Lung cancer is the most common cause for neoplastic LAP
 - Lymphoma
 - Small cell lung cancer can mimic lymphoma
- Neoplastic adenopathy **很少產生 calcification**，除了：
 - Osteogenic sarcoma (OGS)
 - Hodgkin's lymphoma s/p R/T

Inflammatory Adenopathy

- Primary TB
- Fungus: Histoplasmosis
- Viral pneumonia (measles pneumonia)
- Infectious mononucleosis
(rare pulmonary infiltrates, spleen enlargement)
- Sarcoidosis
- AIDS

Inflammatory Adenopathy (1)

■ Primary TB:

■ Primary complex (Ghon complex)

■ Localized **air space consolidation** (esp. in the **middle or lower lobe**) + **LAPs** in a child is a classic CXR presentation.

■ After recovery, LN calcification can occur.

■ Histoplasmosis:

■ In **endemic area**, more common than TB.

■ Tend to have **bulky nodes** and higher incidence of **calcification** than TB.

Inflammatory Adenopathy (2)

■ Viral pneumonia:

- Particularly measles pneumonia.
- Pulmonary infiltrates + mediastinal LAPs in child.
- D.D: primary TB, fungus infection

■ Infectious mononucleosis:

- 常有 widespread LAP, 但是少有 pulmonary infiltrates
- 常有 spleen enlargement

Inflammatory Adenopathy (3)

■ Sarcoidosis:

- Characteristic CXR: 1-2-3 sign

- symmetric hilar LAPs + right paratracheal LAPs

■ AIDS:

- ★ ■ Massive LAPs in AIDS 須考慮:

- Infection: TB, NTM

- Malignancy: NHL, Kaposi's sarcoma

- Drug reactions

Inhalational Disease Adenopathy

- **Silicosis** and **pneumoconiosis** are well-known causes of both hilar and mediastinal LAPs.
- Invariably associated with pulmonary disease: reticulonodular infiltrations.

- **Eggshell calcification**

 ■ 會產生 eggshell calcification 的疾病：

- Silicosis: most common and typical
- Sarcoidosis
- TB
- Lymphoma post R/T

Esophageal Lesions

■ Esophageal cancer

- Abnormal azygoesophageal recess(30%)

- M/3 and L/3 eso. Ca. 可能造成azygoesophageal recess變寬和
azygoesophageal line convex to R't side

- Widened mediastinum

- Tracheal deviation

- Retrocardiac mass

最常見

- Esophageal air-fluid level

- Posterior tracheal indentation or mass

最常見

- Widened posterior tracheal stripe

- Post. tracheal strip: normal - $\leq 3.5\text{mm}$, abnormal - $\geq 4\text{mm}$

- Causes: paratracheal/paraesophageal lymphatic engorgement, retained secretion by distal obstruction by tumor, U/3 tumor itself

Esophageal Lesions

■ Achalasia:

- CXR is usually normal in early phase
- Azygoesophageal recess becomes more convex to the right as the esophagus dilates and is seen as an opacity behind the right border of the heart

■ Operation

- S/p radical resection and reconstruction with an intrathoracic stomach, or colon (usually R't side) pulled up through the mediastinum, retrosternally or subcutaneously.
- Imaging:
 - Widening of mediastinum, usually R't side
 - Air-fluid level when upright
 - Linear shadow representing colon haustra over retrosternal region
 - Subcutaneous and mediastinal emphysema: leak from reconstruction

Post. Mediastinum Lesion

Table 3
Posterior Mediastinal Masses

Esophageal lesions, hiatal hernia
Foregut duplication cyst
Descending aortic aneurysm
Neurogenic tumor
Paraspinal abscess
Lateral meningocele
Extramedullary hematopoiesis

Mediastinal Compartments

Posterior Mediastinal Lesions: V-NAC

1. V: Vertebra

2. N: Nerve-related

3. A: Aorta: D-aorta


Abdomen: Bochdalak's hernia

4. C: Duplication Cyst

● Anterior

● Middle

● Posterior



Vertebra

- **TB spine (paraspinal abscess)**
- **Extramedullary hematopoiesis (EMH)**

Tuberculous Abscess (TB spine)

■ 好發位置:

★ ■ Lower T (thoracic) and upper L (lumbar) vertebrae

■ Early manifestations:

■ Irregularity of the vertebral end plates

★ ■ Decreased height of the intervertebral disc space (吃軟不吃硬)

■ Progressive disease:

■ Anterior wedging of the vertebral body

■ Kyphosis

★ ■ Paravertebral abscess: bulging of paraspinal line

★ ■ Little or no reactive sclerosis or local periosteal reaction

■ Helps distinguish it from pyogenic infections of the spine

★ ■ Rarely affects the posterior vertebral elements (including the pedicles)

■ Distinguish it from metastatic disease

Extramedullary Hematopoiesis (1)

- Common causes of EMH:
 - Thalassemia intermedia or major
 - Congenital hemolytic anemia
 - Congenital spherocytosis
 - Sickle cell anemia
- The most common sites:
 - Liver, spleen
 - Spinal cord: **paravertebral** areas of the thorax
 - Pleura, pulmonary parenchyma, and bronchial wall
- Most common site of intrathoracic sites:
 - **Paravertebral region in lower thorax**

Extramedullary Hematopoiesis (2)

■ CXR:

■ Posterior mediastinal mass

- 單側 or 雙側
- 位置：最常出現在 paravertebral area (along costo-vertebral junction)，lower T-vertebrae (below T6)，甚至可以 involve 整個 vertebrae
- 邊緣清楚、lobulated，以 vertebral body 為中心
- 很少有鈣化 or bone erosion

■ Ribs:

- Marrow expansion 造成 rib 變寬，尤其在 近 vertebra 端
- Trabecula 變得明顯

N: Neurogenic Tumors

- Posterior mediastinal mass 中最常見

- 分成兩類：

- **Nerve root tumor** (schwannoma, neurofibroma): more common in adult, most are benign.

- Schwannoma: The most common neural tumor

- **Sympathetic ganglion tumor** (neuroblastoma, ganglioneuroblastoma, ganglioneuroma): more common in infant and childhood.

Nerve root tumor

- Including schwannoma, neurofibroma, peripheral nerve sheath tumor
- CXR:
 - Round or elliptical (圓形或橢圓形), sharp-marginated mass extending 1-2 rib interface (extent: <4 vertebral body)
 - Often centered at neural foramen; widened neural foramen on lateral view
 - **Cervicothoracic sign**: air-soft tissue interface continues above clavicle indicating posterior location
 - Incomplete border sign

Thank you for Your Attention!